

MIXED LAYER EVOLUTION AND MESOSCALE VARIABILITY IN THE LABRADOR SEA

Dr. Mark Prater and Dr. Tom Rossby
Graduate School of Oceanography
University of Rhode Island
South Ferry Road
Narragansett, RI 02882
mark@seip.gso.uri.edu & tom@rafos.gso.uri.edu
Voice: 401-874-6512 & 401-874-6521, FAX: 401-874-6728
Award No. N000149510509

LONG TERM GOALS

Our long-term goals are to understand the mesoscale dynamics of deep convection, and the links between the surface mixed layer, the underlying thermocline, the upper level circulation, and surface heat fluxes.

OBJECTIVES

We want to examine the mesoscale variability during and immediately following deep convection in the Labrador Sea. We hope to track eddies formed by the baroclinic instability of the chimney and the rim current. We will monitor the evolution of the heat content and the turbulent heat fluxes in the water column before, during, and after deep convection.

APPROACH

We use acoustically tracked, neutrally buoyant RAFOS floats to “tag” and follow water in the Labrador Sea during deep convection events. The floats obtain three navigation fixes per day from four moored sound sources, allowing us to explore Lagrangian pathways and mesoscale variability. During the convection season, between 150-200 temperature (T , to 0.001 °C resolution) and pressure (P , to 0.5 dbar) measurements are collected each day. Knowing the depth response of the float to vertical water velocities (w), we can determine the vertical velocity of the water from the floats’ pressure record. Combining precise temperature data with these velocities allows us to estimate the turbulent heat flux $\langle w'T \rangle$. In addition, we have modified the floats by adding a 30 cm³ volume changing (vocha) mechanism. This vocha is capable of changing the density of the float by about 2 kg m⁻³ at discrete 0.15 kg m⁻³ intervals. We have used this capability by sending the float on repeated profiles of the water column collecting T and P , allowing us to monitor the evolution of the mixed layer depth and heat content. It is important to realize that these measurements are being made in a Lagrangian framework, moving with the water. Difficulties accounting for heat budgets from a fixed, stationary position are thus avoided.

ACCOMPLISHMENTS

The four sound sources used for float navigation and four convection RAFOS floats were deployed from the C.S.S. Hudson in the fall of 1996. During the winter 1997 cruise of the R/V Knorr an additional 18 floats were launched, with 9 more from the spring 1997 cruise of the C.S.S. Hudson. We did, however suffer an unexpectedly large failure rate; out of the 31 floats launched, 14 were not heard from again or returned no useful data. An additional 14 floats

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED 00-00-1997 to 00-00-1997	
4. TITLE AND SUBTITLE Mixed Layer Evolution and Mesoscale Variability in the Labrador Sea				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Rhode Island, Graduate School of Oceanography, Narragansett, RI, 02882				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

surfaced prematurely as a fail-safe response to the float's on-board computer locking-up. After a summer's worth of intensive laboratory and cold room testing, we have pin-pointed the problem and have implemented repairs. Testing and confirmation will continue until the next and final cruise, when the remaining 20 floats will be launched from the upcoming January 1998 cruise of the R/V Knorr.

In spite of our difficulties, we did obtain 60 vertical profiles to almost 1000 m depth, 11 float-months of subsurface Lagrangian trajectories, and over 400 *T/P* time-series. The acoustic navigation is working very well, even in the wind-roughened, surface-trapped sound channel found in the Labrador Sea winter.

SCIENTIFIC/TECHNICAL RESULTS

The RAFOS floats have monitored mixed layer cooling prior to convection (Fig. 1), where the 100 m deep mixed layer cooled by 1 °C in 30 days, indicating a surface heat flux of 150 W m⁻². During the early convection season (mid-February to early March), the mixed layer deepened from 600 m to in excess of 1000 m (the maximum extent of our instrument) (Fig. 2). The change in water column heat content over these 20 days results in a heat flux of 340 W m⁻². At this same time, we also observed downward vertical velocities of nearly near 8 cm s⁻¹. Preliminary correlations with temperature perturbations give turbulent heat fluxes around 150 W m⁻², but more analysis is required to improve these estimates. Finally, floats deployed near Greenland after convection show eddies, both cyclonic and anticyclonic, spinning around the edge of the basin, with one float experiencing over 20 revolutions (Fig. 3). The relative vorticity of this eddy was 0.3 to 0.4 times that of the coriolis parameter *f*.

We have demonstrated the capability of the modified RAFOS float to measure changes of heat content of the mixed layer, and we are able to use the *T* and *P* time-series data to estimate turbulent heat fluxes. We are presently working with surface flux data obtained from Dr. Peter Guest of the Naval Postgraduate School, and will collaborate with him in float/field data intercomparisons.

IMPACT FOR SCIENCE

We consider the profiling RAFOS float to be an effective tool for the monitoring the evolution of the surface waters in a Lagrangian framework. The float, with minimal changes in hardware and software, can also monitor obduction/subduction of thermocline waters before/after the winter cooling season.

TRANSITIONS

No technology transfers or transitions were accomplished this year pertaining to this project.

RELATED PROJECTS

RAFOS floats are currently being deployed as part of the Atlantic Climate Change Experiment (ACCE) in the extension region of the North Atlantic Current (NAC). Our goals in this project are (i) to explore the structure of the boundary between the subpolar and subtropical gyres, and (ii) the sites and mechanisms of exchange. The floats used are isopycnal, and measure temperature, pressure, and oxygen for their nearly two-year missions. One sound source deployed as part of the deep convection experiment in the Labrador Sea provides acoustic navigation in the western subpolar basin. We anticipate some floats deployed in the ACCE

program may be entrained by the Irminger or Greenland Currents and thus map out warm-water pathways into the Labrador Sea.

PUBLICATIONS

Rossby, T. 1997: Numerical experiments with a fluid heated nonuniformly from below. *Tellus*, (resubmitted).

The LABSEA Group, 1997: The Labrador Sea Deep Convection Experiment. *Bull. Amer. Meteor. Soc.* (in preparation).




Figure 1. Temperature profiles from RAFOS float 375, launched in October, 1996. The first six profiles were made at 3 day intervals, the last two profiles were at 5 day intervals. The steady cooling and gradual deepening of the mixed layer can be clearly seen, from which an estimate of a 150 W m^{-2} surface heat flux can be made. The profiles have been offset to one another by $0.5 \text{ }^{\circ}\text{C}$.




Figure 2. Temperature profiles from RAFOS float 388, launched in February, 1997. The profiles were made at 5 day intervals. The mixed layer deepening and concurrent heat loss over the first 20 days of the record can be attributed to a surface heat flux of 340 W m^{-2} . The profiles have been offset to one another by $0.5 \text{ }^{\circ}\text{C}$.




Figure 3. Trajectory of RAFOS float 404, launched off Greenland in May, 1997. The float was tracked 3 times a day, and was trapped in a cyclonic eddy having relative vorticity 0.3 to 0.4 f . The numbers along the float path denote positions of the weekly vertical temperature profiles shown in the bottom panel. The profiles have been offset to one another by 0.5 °C.